

This will help them give you the right medicine at the right time.

- Show this card to your pharmacist
- Show this card to your health care provider

**TO YOU THE PATIENT:**

**for your Pharmacist and Health care Providers**



**TIKOSYN INFORMATION CARD**

To minimize the risk of serious ventricular arrhythmias, patients initiated or re-initiated on TIKOSYN® (dofetilide) should be placed for a minimum of 3 days in a facility that can provide:

- calculations of creatinine clearance
- continuous electrocardiographic monitoring (EKG)
- and cardiac resuscitation

The most common side effects of Tikosyn are headache, chest pain and dizziness.



TK207150 ©2004 Pfizer Inc. All rights reserved. Printed in USA / August 2004

My current dose is \_\_\_\_\_ mcg \_\_\_\_\_ times a day.

- A prescription can only be filled through an enrolled retail pharmacy or a designated mail order pharmacy.
- Tikosyn can be prescribed only by a professional who has completed a Tikosyn education program.

for me.

My doctor has prescribed TIKOSYN® (dofetilide)

**PRESCRIPTION:**

**TO PHARMACIST:**

Tikosyn is only available through the T.I.P.S.<sup>™</sup>  
(Tikosyn In Pharmacy System) program.

**FOR MORE INFORMATION**

please call 1-877-TIKOSYN (1-877-845-6796)  
or go to [www.tikosyn.com](http://www.tikosyn.com)

**1-877-TIKOSYN**

Tikosyn is not for everyone. Please see the enclosed  
full prescribing information and patient product  
information. They contain important information  
about the risk of serious ventricular arrhythmias which  
Tikosyn can cause. The risk can be increased by certain  
drug-drug interactions.

**FOR MORE INFORMATION**

please call 1-877-TIKOSYN (1-877-845-6796)  
or go to [www.tikosyn.com](http://www.tikosyn.com)